

Procedure Flowchart

Design PCR primers based on DNA sequence of reference strain or individual

-

Mix genomic DNAs of the reference and test strains

PCR-amplify DNA regions across the genome, either single reaction or multiplexed reactions


Combining single and/or multiplexed PCR reactions

→

Denature and anneal PCR products to form heteroduplexes/homoduplexes

Scan by temperature gradient electrophoresis to identify heterozygous DNA fragments/variants with a single SNP/simple Indel formed by mixing reference and test strains

Determine sequences of variants to locate the exact positions of the SNP/Indel



Homoduplex heteroduplex formation

omozygote

---GATGAGGCATTTGAC---- CTACTCCGTAAACTG

からこです。

---GATGAGGCATTTGAC-----CTACTCCGTAAACTG

eterozygot

--GATGAGGCATTTGAC--CTACTCCGTAAACTG-


Denature, a

に下る。

-GATGAGGCGTTTGAC --- CTACTCCGCAAACTG

CTACTCCG AAACTG -GATGAGGCGTTTGAC--GATGAGGCATTTGAC--GATGAGGCATTTGAC--CTACTCCGTAAACTG--GATGAGGCGTTTGAC--CTACTCCGCAAACTG-

CTACTCCGTAAACTG



Sample pooling

